

Please carefully read the information of use!

NCL eye drops

Instructions of use

Caution:

- Do not touch the pipette tip of the bottle to any surfaces, as this may lead to contamination of the solution.
- Do not transfer the solution to other bottles or containers.
- Do not use if the solution has changed color.
- If you feel discomfort in the eyes, tearing, change vision, redness of the eyes, stop using the product and consult your doctor because these symptoms can lead to serious diseases.
- Seek medical attention if swallowed.

Precautions :

- Always wash and dry your hands before using eye drops.
 - Close the bottle cap after use.
 - Keep the bottle tightly closed when not in use.
 - Do not store above 30°C.
 - Use before the expiry date specified on the packaging of the product.
 - Do not use more than 60 days after the first opening of the bottle.
 - Do not use if the bottle has been opened or damaged.
 - If more than one topical ophthalmic product is used, the drops should be administered at least fifteen (15) minutes apart.
-

Release form:

Eye drops. 10 ml , in a polyethylene (LDPE) bottle, with a built-in dropper and a polyethylene cap that is screwed on and with the control of first opening. 1 bottle in a pack.

Composition:

Each ml of solution contains : sodium chloride 50 mg .

Excipients: mannitol, boric acid, sodium tetraborate, sodium perborate, sodium hydroxide, hydrochloric acid or sodium hydroxide, water for injections.

Properties:

NCL is a hypertonic ophthalmic solution that removes excess fluid from the cornea due to the osmotic effect. This reduces corneal swelling and associated symptoms.

Sodium perborate, a disappearing preservative, preserves the sterility of the solution in the bottle and disappears when instilled into the eyes, as it decomposes into oxygen and water, so it does not remain in the eyes.

Recommendations for use:

For temporary relief in the case of corneal swelling after eye surgery, after trauma, to relieve swelling caused by inflammation and infection, or secondary to various eye diseases. Irrigation during the provision of first aid when removing foreign particles from the eye.


It is necessary to consult a doctor before use.

Method of application and dosage:

Instill 1 drop into the affected eye every 3-4 hours or as prescribed by the doctor.

How to use NCL eye drops:

Remove the cap from the bottle. Hold the bottle upside down in one hand between the thumb and middle finger. Bring the tip of the bottle to your eye. Do not let it touch your eyes. Dispense one drop into the conjunctival sac formed between your eye and the eyelid by lightly tapping or pressing the base of the vial with your index finger (see picture). Do not squeeze the bottle.

**Adverse reactions:**

For some time after using the drops, the vision may be temporarily blurred.

Rarely, there may be redness, burning, pain, swelling in or around the eyes, which quickly passes. In case of possible occurrence of adverse reactions associated with hypersensitivity to any of the components, consult a doctor.

Use during pregnancy and lactation:

There is no information regarding the safety of use during pregnancy and lactation.

Contraindication:

Increased sensitivity to one of the components.

Storage conditions:

Store at a temperature no higher than 30°C and out of the reach of the children.

Expiration date:

2 years. Use within 60 days after first opening.

Packaging:

Each LDPE bottle contains 10 ml of sterile eye drops.

Manufacturer:

ARISTOPHARMA LTD.

7, Purana Paltan Line, Dhaka-1000, Bangladesh.

Unit 2: Gachha, Gazipur Sadar, Gazipur, Bangladesh.

Name and location of the authorized representative in Ukraine:

LLC “Aristovision”

Kyiv-03022, Ukraine

prov. Kolomyyskyi 17/31a

office-1652

+38 044 257 32 57

e-mail: aristovision.office@gmail.com

The date of the last revision of the instructions for use:

25/03/2024



UA.TR.121

Markings are printed on the packaging:	
	Sterile. Sterilised using aseptic processing methods
	Read the instructions for use
	Batch code (lot)
	Date of manufacture (produced)
	Store at a temperature not exceeding 30°C
	Date of use (use by, month and last digits of the year)
	Manufacturer
	Mark of conformity with technical regulations
	Catalogue number
UA.TR.121	Identification number of the conformity assessment body