

Please carefully read the information of use!

Protear eye drops

Instructions of use

Caution:

- Do not touch the pipette tip of the bottle to any surfaces, as this may lead to contamination of the solution.
- Do not transfer the solution to other bottles or containers.
- Do not use if the solution has changed color.
- If you feel discomfort in the eyes, tearing, change vision, redness of the eyes, stop using the product and consult your doctor because these symptoms can lead to serious diseases.
- Seek medical attention if swallowed.

Precautions :

- Always wash and dry your hands before using eye drops.
- Close the bottle cap after use.
- Keep the bottle tightly closed when not in use.
- Do not store above 30°C.
- Use before the expiry date specified on the packaging of the product.
- Do not use more than 60 days after the first opening of the bottle.
- Do not use if the bottle has been opened or damaged.
- If more than one topical ophthalmic product is used, the drops should be administered at least thirty (30) minutes apart.

Release form:

Eye drops. 10 ml , in a polyethylene (LDPE) bottle, with a built-in dropper and a polyethylene cap that is screwed on and with the control of first opening. 1 bottle in a pack.

Composition:

Each ml contains povidone (colidon 17 PF) 50 mg (mg).

Excipients: sodium chloride, potassium chloride, calcium chloride dihydrate, magnesium chloride, boric acid, sodium lactate solution, sodium perborate, sodium hydroxide or hydrochloric acid, water for injections.

Properties:

Dry eyes can occur when the mucin layer of the tear film is disturbed. The lack of mucin can lead to the destruction of the protective tear film, and as a result to the formation of so-called dry spots on the cornea of the eye, which can cause superficial punctate keratitis in the future. Dry eye syndrome can manifest as burning, itching, photophobia, and a feeling of irritation ("sand") in eyes. "Protear" eye drops contain low molecular pyrogen free povidone with a molecular weight of 7000-11000. Low-molecular-weight povidone (Protear eye drops) has detoxifying properties that can reduce the effect of toxins on the surface of the eye. It is also useful in severe cases of ocular allergic reaction, as it forms a protective layer on the surface of the eye.

After local application, Protear eye drops quickly spread over the conjunctiva and cornea, acting as a substitute for mucin, preventing the development of eye irritation and, as a result, dehydration of the eye epithelium.

At the same time, it forms a moisturizing and protective layer due to its lubricating properties, ability to be absorbed on the surface of the eye, reduce surface tension and increase the viscosity of aqueous solutions. This makes it possible to use it for dry eye syndrome.

Sodium perborate, a disappearing preservative, preserves the sterility of the solution in the bottle and disappears when instilled into the eyes, as it decomposes into oxygen and water, so it does not remain in the eyes.

Recommendations for use:

- With a feeling of burning, itching, photophobia, irritation, dryness and tiredness of the eyes, caused by a decrease in production or absence of the secretion of the lacrimal glands.
- In patients with ocular allergies, Protear is useful as it forms a protective layer that blocks allergens - pollen and dust mites;
- It can be used as a detoxifier .
- It is also sometimes used for corneal ulcers as it covers the cornea with a protective film and therefore helps in the healing process.
- May help contact lens users, those who complains of dry eyes.
- Can be useful after surgery and laser interventions to compensate for the deficiency of the tear film layer. Before use, please consult with doctor.

Method of application and dosage:


Shake the bottle well before use. Adults 1-2 drops in one eye 3-4 times a day. If necessary, it can be used more often.

Before using the drops, contact lenses should be removed and installed not earlier than after 30 minutes after instillation.

When using additional drops, the interval between instillations should be at least 30 minutes. "Protear" should always be the last.

How to use Protear eye drops:

Remove the cap from the bottle. Hold the bottle upside down in one hand between the thumb and middle finger. Bring the tip of the bottle to your eye. Do not let it touch your eyes. Dispense one drop into the conjunctival sac formed between your eye and the eyelid by lightly tapping or pressing the base of the vial with your index finger (see picture). Do not squeeze the bottle.

**Adverse reactions:**

In general, eye drops are well tolerated. In case of possible occurrence of adverse reactions associated with hypersensitivity to any of the components, consult with a doctor.

The ability to influence the reaction speed when driving a motor vehicle or working with other mechanisms: If the drops caused blurred vision, you should wait until the vision clears.

Use during pregnancy and lactation:

There is no information regarding the safety of use during pregnancy and lactation.

Contraindication:

Increased sensitivity to one of the components.

Storage conditions:

Store at a temperature no higher than 30°C and out of the reach of the children.

Expiration date:

2 years. Use within 60 days after first opening.

Packaging:

Each LDPE bottle contains 10 millilitres (ml) of sterile eye drops.

Manufacturer:

ARISTOPHARMA LTD.

7, Purana Paltan Line, Dhaka-1000, Bangladesh.

Unit 2: Gachha, Gazipur Sadar, Gazipur, Bangladesh.

Name and location of the authorized representative in Ukraine:

LLC "Aristovision"

Kyiv-03022, Ukraine

prov. Kolomyyskyi 17/31a

office-1652

+38 044 257 32 57

e-mail: aristovision.office@gmail.com

The date of the last revision of the instructions for use:

25/03/2024



UA.TR.121

Markings are printed on the packaging:	
	Sterile. Sterilised using aseptic processing methods
	Read the instructions for use
	Batch code (lot)
	Date of manufacture (produced)
	Store at a temperature not exceeding 30°C
	Date of use (use by, month and last digits of the year)
	Manufacturer
	Mark of conformity with technical regulations
	Catalogue number
UA.TR.121	Identification number of the conformity assessment body